

10083233

## **IN THE CLAIMS:**

Please amend the claims to read as follows:

- 1. (Currently Amended) A kit for carrying out a nucleic acid amplification reaction, wherein said kit comprises a pair of primers, wherein a at least one primer of said pair contains a modified nucleotide within the three 3' terminal nucleotide positions; wherein said modified nucleotide is selected from the group consisting of 2'-O-methyl nucleotides, 2'-fluoro-nucleotides, 2'-amino nucleotides, and arabinose nucleotides.
- 2. (Original) A kit of claim 1, wherein said modified nucleotide is a 2'-O-methyl nucleotide.
- 3. (Original) A kit of claim 1, wherein said modified nucleotide is a 2'-fluoro-nucleotide.
- 4. (Original) A kit of claim 1, wherein said modified nucleotide is a 2'-amino nucleotide.
- 5. (Original) A kit of claim 1, wherein said modified nucleotide is an arabinose nucleotide.
- 6. (Original) The kit of claim 2, wherein said modified nucleotide is at the 3' terminal position.
- 7. (Original) The kit of claim 3, wherein said modified nucleotide is at the 3' terminal position.
- 8. (Original) The kit of claim 4, wherein said modified nucleotide is at the 3' terminal position.
- 9. (Original) The kit of claim 5, wherein said modified nucleotide is at the 3' terminal position.
- 10. (Original) A kit of claim 1, wherein each primer of said pair of primers independently contains a modified nucleotide within the three 3' terminal nucleotide positions;

wherein said modified nucleotide is selected from the group consisting of 2'-O-methyl nucleotides, 2'-fluoro-nucleotides, 2'-amino nucleotides, and arabinose nucleotides.

- 11. (Currently Amended) A method for amplifying a nucleic acid target sequence, wherein said method comprises carrying out a primer-based amplification reaction in a reaction mixture comprising a pair of primers, wherein a at least one primer of said pair contains a modified nucleotide within the three 3' terminal nucleotide positions; wherein said modified nucleotide is selected from the group consisting of 2'-O-methyl nucleotides, 2'-fluoro-nucleotides, 2'-amino nucleotides, and arabinose nucleotides.
- 12. (Original) The method of claim 11, wherein said modified nucleotide is a 2'-O-methyl nucleotide.
- 13. (Original) The method of claim 11, wherein said modified nucleotide is a 2'-fluoro-nucleotide.
- 14. (Original) The method of claim 11, wherein said modified nucleotide is a 2'-amino nucleotide.
- 15. (Original) The method of claim 11, wherein said modified nucleotide is an arabinose nucleotide.
- 16. (Original) The method claim 12, wherein said modified nucleotide is at the 3' terminal position.
- 17. (Original) The method claim 13, wherein said modified nucleotide is at the 3' terminal position.
- 18. (Original) The method claim 14, wherein said modified nucleotide is at the 3' terminal position.
- 19. (Original) The method claim 15, wherein said modified nucleotide is at the 3' terminal position.

20. (Original) A method of claim 11, wherein each primer of said pair of primers independently contains a modified nucleotide within the three 3' terminal nucleotide positions; wherein said modified nucleotide is selected from the group consisting of 2'-O-methyl nucleotides, 2'-fluoro-nucleotides, 2'-amino nucleotides, and arabinose nucleotides.

-5-